

Customers Information Dos & Don'ts

The PUREBEAU FIBROBLAST handpiece produces at its tip a tiny, so-called plasma flash. This flash leaps over from the tip to the skin and lets it evaporate superficially there in a tiny, micromillimeter-sized point. This superficial point minimizes the skin in its immediate vicinity.

Further points are placed and generate a reduction grid. By the way of their arrangement, these points can equalize skin excesses.

Result: the skin area is markedly sharpened.

How is the treatment?

The treatment begins with the application of an anesthetic cream. For treatment, the PUREBEAU Fibroblast handpiece is used to create small spots in a specific pattern to trigger a tightening reaction.

SECTION

The treatment results in the formation of small, dry spots. The crusts require 5-7 days to fall off and should be covered with makeup. Subsequently slight reddening is visible for 2-3 weeks. No bleeding is to be expected. Blepharoplasty treatment (eyelid tightening) can cause swelling with different symptoms for 1-3 days.

NUMBER OF SESSIONS

In order to achieve an optimal result, several sessions are usually necessary. The easier the treatment intensity, the more sessions are necessary. The number of sessions depends also on the severity of the findings, as well as on the age of the customer.

Normally, 2 to 4 sessions are sufficient, with a minimum time of 6 weeks.

How should you prepare for the treatment?

Tell your beautician before treatment, if you taking any medications or have recently taken (antibiotics, coagulation inhibitors, isotretinoin, etc.) and whether or not you have recently undergone surgery or other aesthetic procedures on your skin.

If you are often suffering from herpes infections, you should also tell them before treatment.

In case of fresh wounds on the face no plasma therapy can be performed.

Otherwise you need to do nothing before the treatment. A healthy lifestyle naturally supports the healing process.

What should you watch after treatment?

Immediately after the treatment, depending on the therapy, small, usually non-bleeding, scab sites appear, which are covered with a special cream after disinfection.

Customers Information Dos & Don'ts

NEVER SCRATCH!

It is important that these crusts are not removed (never scratch, otherwise a scar may remain). Instead, twice daily in the morning and in the evening, carefully clean the areas, then disinfect it with octenisept spray and gently dab with a clean, lint-free cloth. Then apply the special cream.

After the independent peeling of the scab crusts (5th to 8th day), a new, pink skin comes to the fore. This skin must be protected from direct sunlight with a high protection factor of factor 30 for at least 3 months.

Sports, swimming and sauna are only permitted once all the crusts have fallen off and the wound is completely closed.

The after-treatment is the most important component of the treatment and must be followed.

Smokers may experience a significant worsening of the blood flow, which significantly increases the risk of wound healing or slowed healing.

Note that after the crusts are lost, a slight redness may remain in the following weeks to months. It usually takes 2-4 weeks (rarely longer) until the skin has reached its normal color again.

What side effects are possible with the treatment?

Overall, the FIBROBLAST is a particularly low-complication and tissue-strengthening treatment.

After initial redness and scabbing, pink skin emerges.

Rashes of the skin are also rare. Both are usually formed after a few weeks.

Important behavior is a consistent sun protection: during at least 3 months no sunbathing and no solarium, as in some cases to pigment shifts of the skin could occur. During this time, use a sunscreen with a high light protection factor (30 or more).

Very rarely reported hypersensitivity or allergic reactions to local anesthesia, wound healing problems, skin infections, scarring, slight swelling, redness, tingling, the occurrence of herpes simplex (bleeding) and skin depigmentation (whitening).

If one of these undesirable side effects occur, it can usually be corrected during post-treatment.

Please inform us as soon as signs of an infection such as excessive redness, pus, blistering or thick, yellow crust formation or any other undesirable or unexpected side effects occur during the post-treatment period.